# a NARCAN

NDC 69547-383-02

0.1 mL bitwassal stryp per unit for use in de noise only

Rx Only

NARCAN (Indioxone HE)

NASAL SPRAY 4 m

Use NARCAN Nasal Spray for known or suspected opioid overdess in adults and children.

Impertant: For use in the nase only.

Do not remove or test the NARCAN' Nesal Spray until ready to use 

This box contains two (2) 4-mg doses of nationous 
HCI nasal Spray

Two Pack

CHECK PRODUCT EXPIRATION DATE BEFORE USE.

how-to guide zine

NARCAN (naloxone) is used to reverse opioid emergencies by competing with opioids (including but not limited to: prescription pain relievers, heroin, and fentanyl) to bind with the same receptors in the brain. Naloxone can be administered through a nasal spray device or as

The shelf life for naloxone vials is 2-3 years and 4 years for NARCAN nasal spray. Nasal spray devices contain 1 dose.

an injection.

This zine goes over how to administer NARCAN nasal spray for a suspected or known opioid overdose.



**Step.** 1: Check for a suspected overdose and try to wake the person.

Signa of potential opioid overdoses

- unresponsiveness/unconsciousness
   the the second to second to the second to th
  - blue lips or fingernailsslowed or stopped breathing and
  - heartbeat
    - cold/clammy skintiny pupils
    - · limp limbs/body

If the person does not wake up/ continues to be unresponsive, move on to step 2.

### Step. 2: Give 1st dose in the nose.

- Lay the person on their back and tilt their head up
  - and tilt their head up
    Hold the nasal spray with your thumb on the bottom of the
  - thumb on the bottom of the plunger

    Insert the nozzle into either
  - nostril

    Press the plunger firmly to give the 1st dose

Step. 3: Call 911 immediately after giving 1st dose.

In Florida, the "911 Good Samaritan Act" provides immunity for both the person who calls 911 and the person who is experiencing the overdose from being charged, prosecuted, or penalized for possession of a controlled substance. Please check the resources on the back for more information on your state's Good Samaritan laws.

**Step. 4:** Wait 2 - 3 mins after the first dose to give the medicine time to work

- If the person wakes up, move onto step. 5.
- if the person does not wake up, continue to give doses every 2-3 minutes until the person wakes up. (It is safe to keep giving doses)

**Step. 5:** Stay until the ambulance arrives even if the person wakes up

Give another dose if the person becomes very sleepy again.

## Harm. Reduction Saves. Lives

#### **ADDITIONAL RESOURCES:**

https://narcan.com/en/

https://my.clevelandclinic.org/health/diseases/24583-opioid-overdose

#### FREE NALOXONE ACCESS BY STATE:

https://nextdistro.org/naloxone? gad source=1&gbraid=0AAAAACU67k SaAr0dQ3gDS4kJ64byVs9xS

#### **600D. SAMARITAN LAWS. BY. STATE:**

https://www.safeproject.us/goodsamaritan-laws/